

# **LOCH ACHNACLOICH SPECIAL AREA OF CONSERVATION (SAC)**

## **CONSERVATION ADVICE PACKAGE**



## Site Details

Site name:	Loch Achnacloich
Map:	<a href="https://sitelink.nature.scot/site/8293">https://sitelink.nature.scot/site/8293</a>
Location:	Highlands and Islands
Site code:	UK0030189
Area (ha):	20.64
Date designated:	17 March 2005

## Qualifying features

Qualifying feature	SCM assessed condition on this site	SCM visit date	UK overall Conservation Status
Naturally nutrient-rich lakes or lochs which are often dominated by pondweed (Natural eutrophic lakes with <i>Magnopotamion</i> or <i>Hydrocharition</i> -type vegetation). [H3150]	Favourable Maintained	27 July 2010	Unfavourable - Bad

### Notes:

Assessed condition refers to the condition of the SAC feature assessed at a site level as part of NatureScot's [Site Condition Monitoring \(SCM\)](#) programme.

Conservation status is the overall condition of the feature throughout its range within the UK as reported to the European Commission under Article 17 of the Habitats Directive in 2019.

## Overlapping Protected Areas

### [Loch Achnacloich SSSI](#)

## Key factors affecting the qualifying features

### Naturally nutrient-rich lakes or lochs which are often dominated by pondweed

Loch Achnacloich is an example of a lowland, naturally base rich loch (type I in the Duigan (2006) classification scheme). This type of loch is unusual for this part of Scotland as most of the lochs like this are found further south in lowland areas. Loch Achnacloich has relatively high natural productivity and species-richness. Even though Loch Achnacloich is naturally relatively rich in nutrients, lochs of this type are still potentially subject to unnatural over-enrichment.

Loch Achnacloich has a small catchment within a sheltered valley and is fed by the Burn of Achnacloich as well as run-off from surrounding agricultural land which has a mix of arable land and improved pasture with cattle grazing. There are also substantial areas of commercial coniferous forestry in the catchment.

The key management issues include nutrient input leading to over-enrichment (via drainage ditches from surrounding agricultural land as well as the possibility of accidental nutrient enrichment from the slurry pit at Dalnacloich), prevention of colonisation by non-native species and changes to the hydrology or quality of water flowing into the loch and potential changes to water temperature.

Further information about this habitat can be found on the [JNCC website](#).

### **Conservation Objectives for naturally nutrient-rich lakes or lochs which are often dominated by pondweed (natural eutrophic lakes with *Magnopotamion* or *Hydrocharition*-type vegetation)**

#### **1. To ensure that the qualifying feature of Loch Achnacloich SAC is in favourable condition and makes an appropriate contribution to achieving favourable conservation status**

Favourable Conservation Status (FCS) is considered at a European biogeographic level. When determining whether management measures may be required to ensure that the conservation objectives for this site are achieved, the focus should be on maintaining or restoring the contribution that this site makes to FCS.

When carrying out appraisals of plans and projects against these conservation objectives, it is not necessary to understand the status of the feature in other SACs in this biogeographic region. The purpose of the appraisal should be to understand whether the integrity of the site (see objective 2) would be maintained. If this is the case then its contribution to FCS across the Atlantic Biogeographic Region will continue to be met. Further details on how these appraisals should be carried out in relation to maintaining site integrity is provided by objective 2 (including parts a, b, and c). If broader information on the feature is available then it should be used to provide context to the site-based appraisal.

Note that “appropriate” within this part of the conservation objectives is included to indicate that the contribution to FCS varies from site to site and feature to feature.

#### **2. To ensure that the integrity of Loch Achnacloich SAC is maintained by meeting objectives 2a, 2b and 2c for the qualifying feature**

The aim at this SAC is to maintain the naturally nutrient-rich lakes or lochs which are often dominated by pondweed habitat in a favourable condition as a contribution to its wider conservation status. Therefore any impacts to the objectives shown in 2a, 2b or 2c below must not persist so that they prevent the achievement of this overall aim.

When carrying out appraisals of plans or projects the focus should be on maintaining site integrity, specifically by meeting the objectives outlined in 2a, 2b and 2c. If these are met then site integrity will continue to be maintained. Note that not all of these will be relevant for every activity being considered. Any impacts on the objectives shown in 2a, 2b or 2c below must not persist so that they prevent the maintenance of site integrity. Temporary impacts on these objectives resulting from plans or projects can only be permitted where they do not prevent the ability of a feature to recover and there is certainty that the features will be able to quickly recover.

This objective recognises that the qualifying habitat is exposed to a wide range of drivers of change. Some of these are natural and are not a direct result of human influences. Such changes in the habitats’ extent, distribution or condition within the site which are brought about by natural processes, directly or indirectly, are normally considered compatible with

the site's conservation objectives. An exception to this is when the favourable condition of a habitat is dependent on halting or managing natural succession. An assessment of whether a change is natural or anthropogenic, or a combination of both, will need to be looked at on a case by case basis.

#### **2a. Maintain the extent and distribution of the habitat within the site**

The extent of the naturally nutrient-rich lakes or lochs which are often dominated by pondweed habitat has been estimated at 4.7ha. This should be maintained.

The area figure has been taken from the Standard Data Form. There should be no measurable net reduction in the extent of the habitat and its distribution throughout the site should be maintained.

This should include the total surface area, depth of water and type and distribution of loch substrate sediments.

This conservation objective is considered to be met if the conditions to ensure the habitat's long-term existence are in place.

#### **2b. Maintain the structure, function and supporting processes of the habitat**

The structure and function of lochs are strongly influenced by activities within their catchment. Changes in land management or development can affect the integrity of the feature which will manifest itself in changes to the loch.

##### **Physical Attributes**

###### *-Surface Area*

Changes to surface area can indicate pressures on the structure and function of lochs. The surface area of a loch may fluctuate slightly naturally. However changes to surface area and the associated change to depth should be avoided as these can adversely affect the character of the loch, particularly the edge vegetation. Artificial fluctuations to depth found in controlled water bodies such as reservoirs should be avoided as these can adversely affect the vegetation.

Changes to the surface area may also indicate a number of pressures such as abstraction, regulation, construction, excessive sediment deposition and natural succession which may be caused by issues in the catchment.

###### *-Hydrological regime*

The hydrology of the loch should be maintained as this affects both water level fluctuations and annual and within year flushing patterns. Flushing is important as it is strongly related to dilution and removal of nutrients and plankton. Changes to the flushing pattern can be caused by factors similar to those affecting area; abstraction, regulation, construction, excessive sediment deposition and natural succession which may occur in the catchment.

###### *-Loch substrate character*

The type and distribution of sediment particles within the loch should be maintained as this will affect the biology of the loch and the availability of habitats. Changes to the substrate character may also be indicative of changes to the area and hydrological regime. Reduction in area or flushing may affect the substrate character as finer sediments become trapped and there is increased input of leaf-litter from scrub encroachment.

###### *-Natural sediment load*

Accumulation of nutrient-rich sediment may have a strong effect on the water quality and biology of the loch. There should be no increases in sediment loading from land

management practices in the catchment, from the shoreline or short term events such as construction.

Evidence is growing that an increase in storm events associated with climate change may increase the amount of sediment deposited in lochs.

*-Connectivity between the loch and the surrounding area*

While a loch is often perceived as a discrete entity the connections between it and the surrounding area are vital to its functioning as part of a natural system. These natural connections should be maintained by, for example, avoiding hard engineering works on the shoreline or loch bed and anything that impedes the exchange of water either on the surface or with the underlying water table.

**Water Quality**

*-Dissolved Oxygen*

Dissolved oxygen in loch water is vital for respiration of all aquatic animals, including fish, as well as for aquatic plants. An artificially high biomass caused by increased loadings of organic matter or algal blooms should be avoided as this can create a heavy demand which causes low levels of dissolved oxygen. Dissolved oxygen is likely to be lowest in July and August. The target for Loch Achnacloch is dissolved oxygen >6.0mg/l during July and August.

*-pH*

This influences many of the chemical processes in lochs such as the binding of phosphorus. Eutrophic lochs should have pH of 7 to 9. Artificial changes through eutrophication or acidification should be avoided as these can have a significant effect on pH.

*-Nutrients*

In general issues with nutrient levels are inferred from effects upon the vegetation. Phosphorus is one of the main nutrients required for plant growth and there is strong correlation between Total Phosphorus (TP) concentration and phytoplankton biomass. The level of TP in Loch Achnacloch should remain below an annual mean of 50ugP/l.

Nitrogen is the other main nutrient important in loch ecosystems. It is generally less likely to be limiting than phosphorus because of the ability of some organisms to fix Nitrogen from the atmosphere. The target for Loch Achnacloch is that annual mean Total Nitrogen should not exceed 1.5mg/l.

Measurements of Chlorophyll a are used as a proxy for algal growth. Phytoplankton is an important part of the processes of a lake ecosystem affecting light penetration and oxygen demand. Excessive chlorophyll a should be avoided as this is usually associated with nutrient enrichment. Dense floating rafts of filamentous algae (or thick algal coatings on aquatic plants) should also be avoided as these are a negative indicator associated with high nutrient levels.

**2c. Maintain the distribution and viability of typical species of the habitat**

The following species that are typical of naturally nutrient-rich lakes or lochs which are often dominated by pondweed should remain widespread in at Loch Achnacloch

*Callitriche brutia var. hamulata*

*Callitriche hermaphroditica*

*Chara aspera*

*Lemna minor*

*Nuphar pumila*

*Potamogeton berchtoldii*

*Potamogeton crispus*  
*Potamogeton obtusifolius*  
*Potamogeton perfoliatus*  
*Potamogeton pusillus*

The viability of the characteristic species is determined by water quality and other conditions that support the plant community such as water clarity. The distribution and viability of typical species is therefore likely to be linked maintaining natural levels of nutrients in Loch Achnacloch.

Alien species can have direct effects upon the natural plant communities through competition. They may also have more subtle effects as the niche they fill is different and this may directly or indirectly affect the rest of the ecosystem. A list of high impact species has been agreed as part of the Water Framework Directive and none of these should be introduced to Loch Achnacloch.

In addition, the loch provides suitable habitat for mute swan and ducks such as wigeon, tufted duck, teal and mallard. These and other birds may have an impact on the nutrient balance in the loch.

### Conservation Measures

Loch Achnacloch SAC is notified as a Site of Special Scientific Interest and management changes described on the SSSI list of Operations Requiring Consent must have prior consent from SNH (NatureScot)

### Current and recommended management for Naturally nutrient-rich lakes or lochs which are often dominated by pondweed

Issue	Measure	Responsible party
Water abstraction	Ensure timing and volume of abstraction that may be proposed in future is not damaging through discussions with SEPA. Drought plans should adequately address the interests of the site.	SEPA
Afforestation and felling	Ensure that additional forestry is not beyond the carrying capacity of the catchment.	Land Managers Scottish Forestry, Forestry & Land Scotland,
	Planning and implementation of forest harvesting operations should better identify high risk areas. Management should include improved pollution control and blocking of drains.	Scottish Forestry, Forestry & Land Scotland, Forestry owners and managers
	Promote adherence to the Forest and Water Guidelines, and published best practice,	Scottish Forestry, Forestry & Land Scotland, Forestry owners and managers

	during forest restructuring and highlight the need to strictly control fine sediment and other diffuse pollution release into the loch.	
Water quality - enrichment	Implement and maintain monitoring of key water quality parameters.	NatureScot/SEPA
	Any development proposals in the catchment should include appropriate measures to minimise sediment run-off and prevent pollutants from entering the loch.	The Highland Council
	Ensure no adverse impacts from diffuse or point sources.  The width of buffer zones of vegetation between agricultural land and the loch should be increased to reduce man-made nitrogen and phosphorous run-off into the loch from surrounding land. Buffer zones should also be applied to watercourses and drainage directly connected and flowing into Loch Achnacloch.	Land Managers SEPA
Sediment load in the loch from un-afforested land	Avoid land management activities that lead to sediment input into the site either directly or indirectly from elsewhere within the catchment.	Land Managers SEPA The Highland Council
Grazing	Soil erosion should be prevented by ensuring grazing avoids poaching of edges of streams that flow into the loch and of the loch itself. Avoid supplementary feeding of livestock close to the loch as this could concentrate livestock close to the water leading to excess nutrient or sediment entering the loch from poaching, heavy dunging or	Land Managers

	uneaten fodder.	
Development	Ensure any development proposals do not adversely affect the site and should include appropriate measures to minimise sediment run-off and prevent pollutants from entering the loch.	Land Managers The Highland Council NatureScot
Invasive species	Maintain surveillance for invasive species and agree action with NatureScot or SEPA if they are found in Loch Achnacloch.	All
	All anglers and other water users (such as canoeists, wild swimmers or researchers) should follow the Check, Clean, Dry biosecurity procedures to help prevent the spread of invasive species	All
Fishing	Stocking and feeding should be avoided as this can cause enrichment. Introduction of some fish species could also affect the food web in the loch.	Marine Scotland, Cromarty Firth Fishery Board

Contact details:

NatureScot  
Fodderty Way  
Dingwall Business Park  
Dingwall  
IV15 9XB  
Tel: 01463 701 610

Approved on 28 April 2020 by:

**Greg Mudge**  
Principal Advisor  
International Designations

**Graham Neville**  
Area Manager  
Northern Isles & North Highland